

PARTIAL BIBLIOGRAPHY: Water birds as vectors of microbial disease

- Alderiso KA and DeLuca N (1999) Seasonal enumeration of faecal coliform bacteria from the faeces of ring billed gulls (*Larus delawarensis*) and Canada geese (*Branta canadensis*). *Applied and Environmental Microbiology* 65: 5628-5630.
“Results indicate the gull feces contain a greater concentration of fecal coliform bacteria per gram (3.68×10^8) than do goose feces (1.53×10^4),....”
- Benton C, Khan F, Monaghan P, Richards WN and Shedden CB (1983) The contamination of a major water supply by gulls (*Larus* sp.): A study of the problem and remedial actions taken. *Water Res* 17: 789-798.
“Investigations showed a significant correlation between the numbers of gulls roosting and the numbers of *E. coli* present in the water.”
- Brown P (1997) Seabirds' dirty diet spread disease: *New Scientist* 22nd March.
- Butterfield J, Coulson JC, Kearsey SV and Monaghan P (1983) The herring gull *Larus argentatus* as a carrier of *Salmonella*. *J Hygiene Cambridge* 91: 429-436.
- Fayer R, Speer CA and Dubey JP (1997) Avian cryptosporidiosis. In *Cryptosporidium and cryptosporidiosis*, Fayer (Ed.), Boca Raton, CRC Press: 1-33.
- Fenlon DR (1981) Seagulls (*Larus* sp.) as vectors of salmonellae: an investigation into the range of serotypes and numbers of salmonellae in gulls' faeces. *J Hygiene Cambridge* 86: 195-202.
- Fenlon DR (1983) A comparison *Salmonella* serotypes found in the faeces of gulls feeding at a sewage works with serotypes present in the sewage. *J Hygiene Cambridge* 91: 47-52.
- Fennel H, James DB and Morris J (1974) Pollution of a storage reservoir by roosting gulls. *Water Treat and Exam* 23: 5-24.
- Ferns PN and Mudge GP (1999) Abundance, diet and salmonella contamination of gulls feeding at sewage outfalls. *Water Res* 34: 2653-2660.
“The abundance of black-headed gulls and herring gulls was significantly correlated with the volume of sewage discharged.”
- Fricker CR (1984) A note on salmonella excretion in the black headed gull (*Larus ribibundus*) feeding at sewage treatment works. *J Appl Bacteriol* 56(3): 499-502.
- Glunder G, Neumann U, and Braune S (1992) Occurrence of *Campylobacter* spp. in young gulls, duration of *Campylobacter* infection and re-infection by contact. *Zentralbl Veterinarmed* 39(2): 119-122.
- Gould DJ and Fletcher MR (1978) Gull droppings and their effects on water quality. *Water Res* 12: 665-672.
“Daily loads of coliform bacteria indicated that the outputs of the two larger species of gull approximated to values quoted for man.....”
- Graczyk TK, Fayer R, Trout JM, Lewis EJ, Farley CA, Sulaiman I and Lal AA (1998) *Giardia* sp. and infectious *Cryptosporidium parvum* oocysts in the feces of migratory Canada geese (*Branta canadensis*). *Applied and Environ Microbiol* 64: 2736-2738.
- Hatch JJ (1996) Threats to public health from gulls (*Laridae*). *Inter J Environ Health Res.* 6: 5-16.
- Hinshaw WR, McNeil E and Taylor TJ (1994) Avian Salmonellosis, types of salmonella and their relation to public health. *Amer J Hygiene* 40: 264-278.
- Johnston WS, Munro D, Reilly WJ and Sharp JC (1981) An unusual sequel to *Salmonella zanzibar*. *J Hygiene (Lond)* 87(3): 525-528.

“...with seagulls being implicated as the vector transmitting infection from the sewage of a local town on to farmland and an adjacent loch.”

Jones F, Smith P and Watson DC (1977) Pollution of a water supply catchment by breeding Gulls and the potential Environmental Health Implications. *J Inst Water Engrs Sci* 32: 469-482.

Karaguzel A, Kokasal I, Baki A, Fahri U, Gok I and Cirav Z (1993) *Salmonella* and *Shigella* carriage by gulls (*Larus* sp.) on the east Black Sea region of Turkey. *Microbios* 74: 77-80.

Levesque B, Brousseau P, Simard P, Dewailly E, Meisels M, Ransay D and Joly J (1993) Impact of the Ring-billed gull (*Larus delawarensis*) on the Microbiological Quality of Recreational Water. *Appl Environ Microbiol* 59: 1228-1230.

“Gulls can contribute to the bacteriological degradation of recreational water.”

Levesque B, Brousseau P, Bernier F, Dewailly E and Joly J (2000) Study of the bacterial content of ring-billed gull droppings in relation to recreational water quality: *Water Res* 34: 1089-1096.

“...there was little difference in bacterial content as a function of age groups, colony or sampling date.”

Neilsen BB (1960) *Salmonella typhimurium* Carriers in Sea-gulls and Mallards as a possible Source of Infection to Domestic Animals. *Nord Vet-Med* 12: 417-424.

Ortiz NE and Smith GR (1994) Landfill sites, botulism and gulls. *Epidemiol Infect* 112: 385-391.

Pavlassek I (1993) Black headed gull (*Larus-ridibundus* L), a new host of *Cryptosporidium baileyi*. *Veterinari Medicina*, 38: 629-638.

Quessy S and Messier S (1992) Prevalence of *Salmonella*, *Campylobacter*. and *Listeria* in ring-billed gulls (*Larus delawarensis*). *J Wildlife Diseases* 28: 526-531.

Tsubokura M, Matsumoto A, Otsuki K, Animas SB and Sanekata T (1995) Drug resistance and conjugative R plasmids in *Escherichia coli* strains isolated from migratory waterfowl. *J Wildlife Diseases* 31: 352-357.

Weiskel PK, Howes BL and Heufelder GR (1996) Coliform Contamination of a Coastal Embayment: Sources and Transport Pathways. *Env Sci Tech* 30: 1872-1881.

“...it appears that management practices in similar settings should account for migratory waterfowl...”

Whelan CD, Monaghan P, Girdwood RWA and Fricker CR (1988) The significance of wild birds (*Larus* sp.) in the epidemiology of *campylobacter* infections in humans. *Epidemiology and Infection* 101: 259-267.